

CUTTING SEQUENCES AND PALINDROMES

JANE GILMAN AND LINDA KEEN

DEDICATED TO BILL HARVEY ON HIS 65TH BIRTHDAY

ABSTRACT. We give a unified geometric approach to some theorems about primitive elements and palindromes in free groups of rank 2. The geometric treatment gives new proofs of the theorems.

1. INTRODUCTION

In this paper we discuss four older more or less well-known theorems about two generator free groups and a more recent one, an enumerative scheme for primitive words. We describe a geometric technique that ties all of these theorems together and gives new proofs of four of them. This approach and the enumerative scheme will be useful in applications. These applications will be studied elsewhere [5].

The main object here is a two generator free group which we denote by $G = \langle A, B \rangle$.

Definition 1. *A word $W = W(A, B) \in G$ is primitive if there is another word $V = V(A, B) \in G$ such that W and V generate G . V is called a primitive associate of W and the unordered pair W and V is called a pair of primitive associates.*

Definition 2. *A word $W = W(A, B) \in G$ is a palindrome if it reads the same forward and backwards.*

In [3] we found connections between a number of different forms of primitive words and pairs of primitive associates in a two generator free group. These were obtained using both algebra and geometry. The theorems that we discuss, Theorems 2.1, 2.2, 2.3 can be found in [3] and Theorem 2.4 can be found in [11], and the enumeration scheme, along with another proof of Theorem 2.4 can be found in [4].

There are several different geometric objects that can be associated to two generator free groups; among them are the punctured torus, the a three holed sphere and the genus two handlebody. Here we focus on

Rutgers Research Council & Yale University.
PSC-CUNY.

the punctured torus and use “cutting sequences” for simple curves to obtain proofs of Theorems 2.1, 2.2, 2.3 and 2.4.

A similar treatment can be made for the three holed sphere. It was in this setting that we first noticed that the palindromes and products of palindromes were inherent in the geometry by looking at the technique developed in Vidur Malik’s thesis [9] for the three holed sphere representation of two generator groups. The concept of a geometric center of a primitive word was inherent in his work. We thank him for his insight.

2. NOTATION AND DEFINITIONS

In this section we establish the notation and give the definitions needed to state the five theorems and we state them. Note that in stating these theorems in the forms below we are gathering together results from several places into one theorem. Thus, for example, a portion of the statements in theorem 2.1 appears in [7] while another portion appears in [3].

A word $W = W(A, B) \in G$ is an expression $A^{n_1} B^{m_1} A^{n_2} \dots B^{n_r}$ for some set of $2r$ integers $n_1, \dots, n_r, m_1, \dots, n_r$.

The first theorem gives one way to find primitive words. From the form we see how to associate a rational p/q to the primitive word.

Theorem 2.1. *$W = W(A, B)$ in $G = \langle A, B \rangle$ is primitive if and only if, up to cyclic reduction and inverse, it has either the form*

$$(1) \quad A^\epsilon B^{n_1} A^\epsilon B^{n_2} \dots A^\epsilon B^{n_p}$$

where $\epsilon = \pm 1$ and $n_j \geq 1, j < p, |n_j - n_{j+1}| \leq 1$ where j is taken mod p , and, if $\sum_{i=1}^p n_i = q$ and p and q are relatively prime, $(p, q) = 1$.

or the form

$$(2) \quad B^\epsilon A^{n_1} B^\epsilon A^{n_2} \dots B^\epsilon A^{n_q}$$

where $\epsilon = \pm 1$ and $n_j \geq 1, j < q, |n_j - n_{j+1}| \leq 1$ where j is taken mod p , and, if $\sum_{i=1}^q n_i = p$ and p and q are relatively prime, $(p, q) = 1$.

In the first case, we denote the primitive word by $W_{p/q}$ and in the second by $W_{q/p}$. Two primitive words $W_{p/q}$ and $W_{r/s}$ are associates if and only $|ps - qr| = 1$.

2.1. Farey arithmetic. In what follows when we use r/s to denote a rational, we assume that r and s are integers, $s \neq 0$ and $(r, s) = 1$. We also use the notation $1/0$ to denote the point at infinity.

To state the second theorem, we need the concept of Farey addition for fractions.

Definition 3. If $\frac{p}{q}, \frac{r}{s} \in \mathbb{Q}$, the Farey sum is

$$\frac{p}{q} \oplus \frac{r}{s} = \frac{p+r}{q+s}$$

Two fractions are called Farey neighbors or simply called neighbors if $|ps - qr| = 1$.

Note that the Farey neighbors of $1/0$ are the rationals $n/1$. If $\frac{p}{q} < \frac{r}{s}$ then it is a simple computation to see that

$$\frac{p}{q} < \frac{p}{q} \oplus \frac{r}{s} < \frac{r}{s}$$

and that both pairs of fractions

$$\left(\frac{p}{q}, \frac{p}{q} \oplus \frac{r}{s}\right) \text{ and } \left(\frac{p}{q} \oplus \frac{r}{s}, \frac{r}{s}\right)$$

are neighbors if $(p/q, r/s)$ are.

We can create the diagram for the Farey tree by marking each fraction by a point on the real line and joining each pair of neighbors by a semi-circle orthogonal to the real line in the upper half plane. The points $n/1$ are joined to their neighbor $1/0$ by vertical lines. The important thing to note here is that because of the properties above none of the semi-circles or lines intersect in the upper half plane. Each pair of neighbors together with their Farey sum form the vertices of a curvilinear or hyperbolic triangle and the interiors of two such triangles are disjoint. Together the set of these triangles forms a tessellation of the hyperbolic plane which is known as the Farey tree.

Fix any point ζ on the positive imaginary axis. Given a fraction, $\frac{p}{q}$, there is a hyperbolic geodesic γ from ζ to $\frac{p}{q}$ that intersects a minimal number of these triangles.

Definition 4. The Farey level or the level of p/q , $Lev(p/q)$ is the number of triangles traversed by γ

Note that the curve (line) γ joining ζ to either $0/1$ or $1/0$ does not cross any triangle so these rationals have level 0. The geodesic joining ζ to $1/1$ intersects only the triangle with vertices $1/0, 0/1$ and $1/1$ so the level of $1/1$ is 1. Similarly the level of $n/1$ is n .

Definition 5. We determine a Farey sequence for $\frac{p}{q}$ by choosing the new endpoint of each new common edge in the sequence of triangles traversed by γ .

Given p/q , we can find the smallest and largest rationals m/n and r/s that are its neighbors. These also have the property that they are the only neighbors with lower level. That is, as rational numbers $m/n < p/q < r/s$ and the levels satisfy $Lev(m/n) < Lev(p/q)$, $Lev(r/s) < Lev(p/q)$, and if u/v is any other neighbor $Lev(u/v) > Lev(p/q)$.

Definition 6. We call the smallest and the largest neighbors of the rational p/q the distinguished neighbors of p/q .

Note that we can tell whether which distinguished neighbor r/s is smaller (respectively larger) than p/q by the sign of $rq - ps$.

Farey sequences are related to continued fraction expansions of fractions (see for example, [6]). In particular, write

$$\frac{p}{q} = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots + \frac{1}{a_k}}}} = [a_0, \dots, a_k]$$

where $a_j > 0$, $j = 1 \dots k$ and set $\frac{p_n}{q_n} = [a_0, \dots, a_n]$. The approximating fractions can be computed recursively from the continued fraction for p/q as follows:

$$p_0 = a_0, q_0 = 1 \text{ and } p_1 = a_0a_1 + 1, q_1 = a_1$$

$$p_j = a_j p_{j-1} + p_{j-2}, q_j = a_j q_{j-1} + q_{j-2} \quad j = 2, \dots, k.$$

The level of p/q can be expressed in terms of the continued fraction expansion by the formula

$$Lev(p/q) = \sum_{j=0}^k a_j.$$

The distinguished neighbors of p/q have continued fractions

$$[a_0, \dots, a_{k-1}] \text{ and } [a_0, \dots, a_{k-1}, a_k - 1].$$

The Farey sequence contains the approximating fractions as a subsequence. The points of the Farey sequence between $\frac{p_j}{q_j}$ and $\frac{p_{j+1}}{q_{j+1}}$ have continued fraction expansions

$$[a_0, a_1, \dots, a_j + 1], [a_0, a_1, \dots, a_j + 2], \dots, [a_0, a_1, \dots, a_j + a_{j+1} - 1].$$

The approximating fractions $\frac{p_j}{q_j}$ are alternately larger and smaller than $\frac{p}{q}$. The number a_j counts the number of times the new endpoint in the Farey sequence lies on one side of the old one.

2.2. Farey words, continued fraction expansions and algorithmic words. The next theorem gives a recursive enumeration scheme for primitive words using Farey sequences of rationals.

Theorem 2.2. *The primitive words in $G = \langle A, B \rangle$ can be enumerated inductively by using Farey sequences as follows: set*

$$W_{0/1} = A, \quad W_{1/0} = B.$$

Given p/q , consider its Farey sequence. Let $\frac{m}{n}$ and $\frac{r}{s}$ be its distinguished neighbors labeled so that

$$\frac{m}{n} < \frac{p}{q} < \frac{r}{s}.$$

Then

$$W_{\frac{p}{q}} = W_{\frac{m}{n} \oplus \frac{r}{s}} = W_{r/s} \cdot W_{m/n}.$$

A pair $W_{p/q}, W_{r/s}$ is a pair of primitive associates if and only if $\frac{p}{q}, \frac{r}{s}$ are neighbors, that is, $|ps - qr| = 1$.

We use the same notation for these words as those in Theorem 2.1 because, as we will see when we give the proofs of the theorems, we get the same words.

We note that the two products $W_{m/n} \cdot W_{r/s}$ and $W_{r/s} \cdot W_{m/n}$ are always conjugate in G . In the iteration scheme we always choose the product where the larger index comes first. The point is that in order for the scheme to work the choice has to be made consistently. We emphasize that $W_{p/q}$ always denotes the word obtained using this enumeration scheme.

The $W_{p/q}$ words can be expanded using their continued fraction exponents instead of their primitive exponents. This is also known as the *algorithmic form* of the primitive words, that is, the form in which the words arise in the $PSL(2, \mathbb{R})$ discreteness algorithm [2, 3]. The following theorem exhibits the primitive words with the continued fraction expansion exponents in its most precise form.

Theorem 2.3. *If $[a_0, \dots, a_k]$ is the continued fraction expansion of p/q , the primitive word $W_{p/q}$ can be written inductively using the continued fraction approximants $p_j/q_j = [a_0, \dots, a_j]$. They are alternately larger and smaller than p/q .*

Set

$$W_{0/1} = A, \quad W_{1/0} = B \quad \text{and} \quad W_{1/1} = BA.$$

If $p_{k-2}/q_{k-2} > p/q$ set

$$W_{p/q} = W_{p_{k-2}/q_{k-2}} (W_{p_{k-1}/q_{k-1}})^{a_k}$$

and set

$$W_{p/q} = (W_{p_{k-1}/q_{k-1}})^{a_k} W_{p_{k-2}/q_{k-2}}$$

otherwise.

The following theorem is proved in [11].

Theorem 2.4. *Let $G = \langle A, B \rangle$ be a two generator free group. Then any primitive element $W \in G$ is conjugate to a cyclic permutation of either a palindrome in A, B or a product of two palindromes. In particular, if the length of W is $p + q$, then, up to cyclic permutation, W is a palindrome if and only if $p + q$ is odd and is a product of two palindromes otherwise.*

We note that this can be formulated equivalently using the parity of pq which is what we do below.

In the pq odd case, the two palindromes in the product can be chosen in various ways. We will make a particular choice in the next theorem.

2.3. E-Enumeration. The next theorem, proved in [4], gives yet another enumeration scheme for primitive words, again using Farey sequences. The new scheme to enumerate primitive elements is useful in applications, especially geometric applications. These applications will be studied elsewhere [5]. Because the words we obtain are cyclic permutations of the words $W_{p/q}$, we use a different notation for them; we denote them as $E_{p/q}$.

Theorem 2.5. *The primitive elements of a two generator free group can be enumerated recursively using their Farey sequences as follows. Set*

$$E_{0/1} = A, E_{1/0} = B, \text{ and } E_{1/1} = BA.$$

Given p/q with distinguished neighbors $m/n, r/s$ such that $m/n < r/s$,

- *if pq is odd, set $E_{p/q} = E_{r/s}E_{m/n}$ and*
- *if pq is even, set $E_{p/q} = E_{m/n}E_{r/s}$. In this case $E_{p/q}$ is the unique palindrome cyclicly conjugate to $W_{p/q}$.*

$E_{p/q}$ and $E_{p'/q'}$ are primitive associates if and only if $pq' - qp' = \pm 1$.

Note that when pq is odd, the order of multiplication is the same as in the enumeration scheme for $W_{p/q}$ but when pq is even, it is reversed. This theorem says that if pq is even, there is a unique palindrome cyclicly conjugate to $W_{p/q}$. If pq is odd, it determines a canonical factorization of $W_{p/q}$ into a pair of palindromes. This factorization exhibits the Farey sequence of p/q and the order of multiplication is what makes the enumeration scheme work.

In this new enumeration scheme, Farey neighbors again correspond to primitive pairs but the elements of the pair $(W_{p/q}, W_{p'/q'})$ are not necessarily conjugate to the elements of the pair $(E_{p/q}, E_{p'/q'})$ by the same element of the group. .

3. CUTTING SEQUENCES

We represent G as the fundamental group of a punctured torus and use the technique of *cutting sequences* developed by Series (see [12, 7, 10]) as the unifying theme. This representation assumes that the group G is a discrete free group. Cutting sequences are a variant on Nielsen boundary development sequences [10]. In this section we outline the steps to define cutting sequences.

- It is standard that $G = \langle A, B \rangle$ is isomorphic to the fundamental group of a punctured torus S . Each element of G corresponds to a free homotopy class of curves on S . The primitive elements are classes of simple curves that do not simply go around the puncture. Primitive pairs are classes of simple closed curves with a single intersection point.
- Let \mathcal{L} be the lattice of points in \mathbb{C} of the form $m + ni$, $m, n \in \mathbb{Z}$ and let \mathcal{T} be the corresponding lattice group generated by $a = z \mapsto z + 1, b = z \mapsto z + i$. The (unpunctured) torus is homeomorphic to the quotient $\mathbb{T} = \mathbb{C}/\mathcal{T}$. The horizontal lines map to longitudes and the vertical lines to meridians on \mathbb{T} .

The punctured torus is homeomorphic to the quotient of the plane punctured at the lattice, $(\mathbb{C} \setminus \mathcal{L})/\mathcal{T}$. Any curve in \mathbb{C} whose endpoints are identified by the commutator $aba^{-1}b^{-1}$ goes around a puncture and is no longer homotopically trivial.

- The simple closed curves on \mathbb{T} are exactly the projections of lines joining pairs of lattice points (or lines parallel to them). These are lines $L_{q/p}$ of rational slope q/p . The projection $l_{q/p}$ consists of p longitudinal loops q meridional loops. We assume that p and q are relatively prime; otherwise the curve has multiplicity equal to the common factor.

For the punctured torus, any line of rational slope, not passing through the punctures projects to a simple closed curve and any simple closed curve, not enclosing the puncture, lifts to a curve freely homotopic to a line of rational slope.

- Note that, in either case, if we try to draw the projection of $L_{q/p}$ as a simple curve, the order in which we traverse the loops on \mathbb{T} (or S) matters. In fact there is, up to cyclic permutation

and reversal, only one way to draw the curve. We will find this way using cutting sequences. Below, we assume we are working on \mathbb{T} .

- Choose as fundamental domain (for S or \mathbb{T}) the square D with corners (puncture points) $\{0, 1, 1+i, i\}$. Label the bottom side A and the top side \bar{A} ; label the left side B and the right side \bar{B} . It will be clear why we use the same letters for the labels as for the generators of G . Use the translation group to label the sides of all copies of D in the plane.
- Choose a fundamental segment of the line $L_{q/p}$ and pick one of its endpoints as initial point. It passes through $p + |q|$ copies of the fundamental domain. Call the segment in each copy a strand.

Because the curve is simple, there will either be “vertical” strands joining the sides A and \bar{A} , or “horizontal” strands joining the sides B and \bar{B} , but not both.

Call the segments joining a horizontal and vertical side corner strands. There are four possible types of corner strands: from left to bottom, from left to top, from bottom to right, from top to right. If all four types were to occur, the projected curve would be trivial on \mathbb{T} . There cannot be only one or three different types of corner strands because the curve would not close up. Therefore the only corner strands occur on one pair of opposite corners and there are an equal number on each corner.

- Traversing the fundamental segment from its initial point, the line goes through or “cuts” sides of copies of D . We will use the side labeling to define a *cutting sequence* for the segment. Since each side belongs to two copies it has two labels. We have to pick one of these labels in a consistent way. As the segment passes through, there is the label from the copy it leaves and the label from the copy it enters. We always choose the label from the copy it enters. Note that the cyclic permutation depends on the starting point.
- If $|q|/p < 1$, the resulting cutting sequence will contain p A 's (or p \bar{A} 's), $|q|$ B 's (or $|q|$ \bar{B} 's) and there will be $p - |q|$ horizontal strands and p corner strands; if $|q|/p > 1$, the resulting cutting sequence will contain p A 's (or p \bar{A} 's), $|q|$ B 's (or $|q|$ \bar{B} 's) and there will be $|q| - p$ vertical strands and $|q|$ corner strands. We identify the cutting sequence with the word in W interpreting the labels A, B and the generators and the labels \bar{A}, \bar{B} as their inverses.

- Given an arbitrary word $W = A^{m_1} B^{n_1} A^{m_2} B^{n_2} \dots A^{m_p} B^{n_p}$ in G , we can form a cutting sequence for it by drawing strands from the word through successive copies of D . Consider translates of the resulting curve by elements of the lattice group. If they are all disjoint up to homotopy, the word is primitive.

Let us illustrate with three examples. In the first two, we draw the cutting sequences for the fractions $\frac{q}{p} = \frac{1}{1}$ and $\frac{q}{p} = \frac{3}{2}$. In the third, we construct the cutting sequence for the word $A^3 B^2$.

- A fundamental segment of $l_{1/1}$ can be chosen to begin at a point on the left (B) side and pass through D and the adjacent copy above D ; There will be a single corner strand connecting the B side to an \bar{A} side and another connecting an A side to an \bar{B} side.

To read off the cutting sequence begin with the point on B and write B . Then as we enter the next (and last) copy of D we have an A side. The word is thus BA .

Had we started on the bottom, we would have obtained the word AB .

- A fundamental segment of $L_{3/2}$ passes through 5 copies of the fundamental domain. There is one “vertical” segment joining an A and an \bar{A} , 2 corner segments joining a B and an \bar{A} and two joining the opposite corners. Start on the left side. Then, depending on where on this side we begin we obtain the word $BABAA$ or $BAABA$.

If we start on the bottom so that the vertical side is in the last copy we encounter we get $ABABA$.

- To see that the word $AAABB$ cannot correspond to a simple loop, draw the a vertical line of length 3 and join it to a horizontal line of length 2. Translate it one to the right and one up. Clearly the translate intersects the curve and projects to a self-intersection on the torus. This will happen whenever the horizontal segments are not separated by a vertical segment.

Another way to see this is to try to draw a curve with 3 meridian loops and two longitudinal loops on the torus. You will easily find that if you try to connect them arbitrarily the strands will cross on \mathbb{T} , but if you use the order given by the cutting sequence they will not. Start in the middle of the single vertical strand and enter a letter every time you come to the beginning of a new strand. We get $ABABA$.

- Suppose $W = A^3 B^2$. To draw the cutting sequence, begin on the bottom of the square and, since the next letter is A again,

draw a vertical strand to a point on the top and a bit to the right. Next, since we have a third A , in the copy above D draw another vertical strand to the top and again go a bit to the right. Now the fourth letter is a B so we draw a corner strand to the right. Since we have another B we need to draw a horizontal strand. We close up the curve with a last corner strand from the left to the top.

Because we have both horizontal and vertical strands, the curve is not simple and the word is not primitive.

4. PROOFS

Proof of theorem 2.1. Suppose first that $0 < p/q \leq 1$. The cases $p/q = 0/1, 1/0$ are trivial. The other cases follow in the same way, either interchanging A and B or replacing B by \bar{B} .

The line $L_{q/p}$ has slope at least 1 so there will be at least one vertical strand and no horizontal strands. Choose as starting point the lowest point on a B side. Because there are no horizontal strands, we must either go up or down; assume we go up. The first letter in the cutting sequence is B and since the strand must be a corner strand, the next letter is A . As we form the cutting sequence we see that because there are no horizontal strands, no B can be followed by another B . Because we started at the lowest point on B , the last strand must start at the rightmost point on an A side. Since there are $p+q$ strands, this means the sequence, and hence the word has the form of equation (1) with $\epsilon = 1$,

$$W_{p/q} = BA^{n_0}BA^{n_2}B \dots A^{n_p}, \quad \sum n_i = q.$$

If we use the translation group to put all the strands into one fundamental domain, the endpoints of the strands on the sides are ordered. We see that if we are at a point on the A side, the next time we come to the A side we are at a point that is p to the right mod(q). The condition $|n_i - n_{i+1}| \leq 1$, follows since we cycle around in this way. This proves Theorem 2.1 in this case.

Note that had we chosen a different starting point we would have obtained a cyclic permutation of $W_{p/q}$, or, depending on the direction, its inverse.

For $1 < q/p$ we have no vertical strands and we interchange the roles of A and B . We obtain a primitive word in the form of equation (2) with $\epsilon = 1$.

For $p/q < 0$, we replace A or B by \bar{A} or \bar{B} as appropriate and obtain the either equation (1) or equation (2) with $\epsilon = -1$.

To see when two primitive words $W_{p/q}$ and $W_{r/s}$ are associates, note that the lattice \mathcal{L} is generated by fundamental segments of lines $L_{p/q}, L_{r/s}$ if and only if $|ps - qr| = 1$, or equivalently, if and only if $(p/q, r/s)$ are neighbors. \square

Proof of theorem 2.2 and 2.3.

The theorem prescribes a recursive definition of a primitive word associated to a rational p/q . We need to show that the products formed in the recursion are the primitive words we obtained in Theorem 2.1. We proceed by induction on the level of p/q .

At level 1, the distinguished neighbors of $1/1$ are $0/1$ and $1/0$. The recursion gives $W_{0/1} = A, W_{1/0} = B$ and $W_{1/1} = BA$ which is a primitive of the correct form.

Note that we have taken the product in the recursion so that the word corresponding to the bigger fraction is on the left. This choice is consistent with the choice we made in defining the cutting sequences.

Assume now that for all fractions with level less than N , the words we obtain are primitive words in the appropriate form and let p/q have level N ; that is, $Lev(p/q) = N$. Assume for argument's sake that $0 < p/q \leq 1$. The other cases follow in a similar way.

Let m/n and r/s be the distinguished neighbors of p/q labeled so that $\frac{m}{n} \oplus \frac{r}{s} = \frac{p}{q}$ and $nr - ms = 1$.

Since these are distinguished neighbors $Lev(m/n) < N, Lev(r/s) < N$ so the words $W(m/n)$ and $W(r/s)$ are primitive and form a pair of primitive associates.

We need to consider the product

$$(3) \quad W_{r/s}W_{m/n} = B^\epsilon A^{k_1} B^\epsilon A^{k_2} \dots B^\epsilon A^{k_r} B^\epsilon A^{l_1} B^\epsilon A^{l_2} \dots B^\epsilon A^{l_m}.$$

By our assumptions on $p/q, 0 \leq m/n < r/s \leq 1$ so the exponents of B are $+1$ in both words. This also implies that $k_r \geq 1, l_m \geq 1$. The conditions $\sum_{i=1}^r k_i + \sum_{i=1}^m l_i = s + n$, and $(m + r, n + s) = 1$ clearly hold for these exponents.

The only condition we have left to check is that $|k_r - l_1| \leq 1$. Assume first $Lev(r/s) < Lev(m/n)$. This implies that r/s is the right distinguished neighbor of m/n and that its left distinguished neighbor u/v also satisfies $Lev(u/v) < Lev(m/n)$.

In terms of continued fractions we see this by writing $p/q = [0, a_1, \dots, a_k]$. $Lev(r/s) < Lev(m/n)$ implies

$$m/n = [0, a_1, \dots, a_{k-1}, a_k - 1]$$

$$r/s = [0, a_1, \dots, a_{k-1}]$$

and

$$u/v = [0, a_1, \dots, a_{k-1} - 1].$$

By the induction hypothesis we have

$$W_{m/n} = W_{r/s}W_{u/v}$$

so that

$$W_{p/q} = W_{r/s}W_{m/n} = W_{r/s}^2W_{u/v}.$$

Writing out the words we have

$$W_{r/s} = BA^{k_1}BA^{k_2} \dots BA^{k_r}$$

$$W_{u/v} = BA^{j_1}BA^{j_2} \dots BA^{j_u}$$

so that

$$W_{m/n} = BA^{l_1}BA^{l_2} \dots BA^{l_m} = BA^{k_1}BA^{k_2} \dots BA^{k_r}BA^{j_1}BA^{j_2} \dots BA^{j_u}.$$

From these expressions we see that $k_1 = l_1$. Now we know that $|k_r - k_1| \leq 1$ because this condition is computed mod r , and therefore $|k_r - l_1| \leq 1$ and $W_{p/q}$ is a primitive word.

The same argument with right and left reversed holds if $Lev(m/n) < Lev(r/s)$.

To complete the proof we note that this also shows if $r/s, m/n$ are neighbors, the pair $W_{r/s}, W_{m/n}$ is a pair of primitive associates. If $r/s, m/n$ are not neighbors, the vectors joining zero with $m + ni$ and $r + si$ do not generate the lattice \mathcal{L} and the classes corresponding to $W_{r/s}, W_{m/n}$ do not generate $\pi_1(S)$ so the words are not primitive associates.

We note that proving Theorem 2.3 is just a matter of notation. □

As a consequence of this theorem, if given a primitive associate pair $(W_{p/q}, W_{r/s})$, we draw the strands for cutting sequence for each primitive in the same diagram, then the result is the cutting sequence of the product.

Proof of Theorem 2.4.

Suppose pq is even. Again we prove the theorem for $0 < p/q < 1$. The other cases follow as above by interchanging the roles of A and B or replacing B by \overline{B} . The idea is to choose the starting point correctly.

Draw a line of slope q/p . By assumption, there are vertical but no horizontal strands and $q - p > 0$ must be odd. This implies that in a fundamental segment there are an odd number of vertical strands. In particular, if we pull all the strands of a fundamental segment into one copy of D , one of the vertical strands is the middle strand. Choose the fundamental segment for the line in the lattice so that it is centered about this middle vertical strand.

To form the cutting sequence for the corresponding word W , begin at the top of the middle strand and move upwards. Now go downwards and form the cutting sequence for a word V . By the symmetry, since we began with a middle strand, V is W with all the A 's replaced by

\bar{A} 's and all the B 's replaced by \bar{B} 's. Since $V = W^{-1}$, we see that W must be a palindrome which we denote as $W = P_{p/q}$. Moreover, since it is the cutting sequence of a fundamental segment of the line of slope q/p , it must be a cyclic permutation of $W_{p/q}$.

Note that since we began by going up along a vertical strand, the first letter in the sequence is an A and, since it is a palindrome, so is the last letter.

When $q/p > 1$, there are horizontal and no vertical strands, and there is a middle horizontal strand. This time we choose this strand and go right and left to see that we get a palindrome. The first and last letters in this palindrome will be B .

If $p/q < 0$, we argue as above but either A or B is replaced by respectively \bar{A} or \bar{B} . \square

We now turn to the enumeration scheme:

Enumeration for Theorem 2.5.

The proof of the enumeration theorem involves purely algebraic manipulations and can be found in [4]. We do not reproduce it here but rather give a heuristic geometric idea of the enumeration and the connection with palindromes that comes from the $PSL(2, \mathbb{R})$ discreteness algorithm [1, 2].

Note that the absolute value of the trace of an element $X \in PSL(2, \mathbb{R})$, $|\text{trace}(X)|$, is well-defined. Recall that X is elliptic if $|\text{trace}(X)| < 2$ and hyperbolic if $|\text{trace}(X)| > 2$. As an isometry of the upper half plane, each hyperbolic element has an invariant geodesic called its *axis*. Each point on the axis is moved a distance $l(X)$ towards one endpoint on the boundary. This endpoint is called the attractor and the distance can be computed from the trace by the formula $\cosh \frac{l(X)}{2} = \frac{1}{2}|\text{trace}(X)|$. The other endpoint of the axis is a repeller.

For convenience we use the unit disk model and consider elements of $PSL(2, \mathbb{R})$ as isometries of the unit disk. In the algorithm one begins with a representation of the group where the generators A and B are (appropriately ordered) hyperbolic isometries of the unit disk. The algorithm applies to any non-elementary representation of the group where the representation is not assumed to be free or discrete. The axes of A and B may be disjoint or intersect. We illustrate the geometric idea using intersecting axes.

If the axes of A and B intersect, they intersect in unique point p . In this case one does not need an algorithm to determine discreteness or non-discreteness as long as the multiplicative commutator, $ABA^{-1}B^{-1}$,

is not an elliptic isometry. However, the geometric steps used in determining discreteness or non-discreteness in the case of an elliptic commutator still make sense. We think of the representation as being that of a punctured torus group when the group is discrete and free.

Normalize at the outset so that the translation length of A is smaller than the translation length of B , the axis of A is the geodesic joining -1 and 1 with attracting fixed point 1 and the axis of B is the line joining $e^{i\theta}$ and $-e^{i\theta}$. This makes the point p the origin. Replacing B by its inverse if necessary, we may assume the attracting fixed point of B is $e^{i\theta}$ and $-\pi/2 < \theta \leq \pi/2$.

The geometric property of the palindromic words is that their axes all pass through the origin.

Suppose $(p/q, p'/q')$ is a pair of neighbors with pq and $p'q'$ even and $p/q < p'/q'$. The word $W_{r/s} = W_{p'/q'}W_{p/q}$ is not a palindrome or conjugate to a palindrome. Since it is a primitive associate of both $W_{p'/q'}$ and $W_{p/q}$ the axis of $Ax_{W_{r/s}}$ intersects each of the axes $Ax_{W_{p/q}}$ and $Ax_{W_{p'/q'}}$ in a unique point; denote these points by $q_{p/q}$ and $q_{p'/q'}$ respectively. Thus, to each triangle, $(p/q, r/s, p'/q')$ we obtain a triangle in the disk with vertices $(0, q_{p/q}, q_{p'/q'})$.

The algorithm provides a method of choosing a next neighbor and next associate primitive pair so that at each step the longest side of the triangle is replaced by a shorter side. The procedure stops when the sides are as short as possible. Of course, it requires proof to see that this procedure will stop and thus will actually give an algorithm.

There is a similar geometric description of the algorithm and palindromes in the case of disjoint axes.

REFERENCES

1. Gilman, Jane *Alternate Discreteness tests* Birman Conf. Proceedings, AMS/IP Studies in Advanced Mathematics, Volume 24, (2001), 41-47.
2. Gilman, Jane *Two-generator Discrete Subgroups of $PSL(2, \mathbf{R})$* , Memoirs of the AMS, volume 561 (1995).
3. Gilman, Jane and Keen, Linda *Word Sequence and Intersection Numbers*, Cont. Math **311** (2002), pp. 331-349.
4. Gilman, Jane and Keen, Linda *Enumerating Plaindromes in Rank Two Free Groups*, submitted.
5. Gilman, Jane and Keen, Linda *Palindromes and discreteness for Kleinian groups*, in preparation.
6. Hardy, G.H. and Wright, E. M., *An introduction to the theory of numbers* Oxford, Clarendon Press 1938.
7. Keen, Linda and Series, Caroline, *Pleating Coordinates for the Maskit Embedding of Teichmüller space for a punctured torus*, Topology, Vol. **32** #4, (1993), 719-749.

8. Magnus, Wilhelm; Karass, Abraham; and Solitar, Donald *Combinatorial Group Theory* (1966) John Wiley & Sons, NYC.
9. Malik, Vidur *Curves Generated on Surfaces by the Gilman-Maskit Algorithm*, Ph D thesis, Rutgers University, Newark, NJ (2007).
10. Nielsen, Jacob *Untersuchungen zur Topologie der geschlossenen zweiseitigen Flächen, I, II III* Acta Math. 50 (1927) 189-358; 53 (1929) 1-76, 58 (1932), 87-167. *Investigation in the Topology of Closed Oriented Surfaces*
11. Piggott, Adam *Palindromic primitives and palindromic bases in the free group of rank two* J. Algebra 304 (2006), no. 1, 359-366.
12. Series, Caroline *The Geometry of Markoff Numbers*, Math. Intelligencer 7 #3,(1985), 20-29.
13. Wright, David J., *Searching for the cusp*, Spaces of Kleinian Groups, LMS Lecture Notes 329, Cambridge U. Press (2004), 1-36.

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NEWARK, NJ 07079
E-mail address: `gilman@andromeda.rutgers.edu`

DEPARTMENT OF MATHEMATICS, LEHMAN COLLEGE AND GRADUATE CENTER, CUNY, BRONX, 10468
E-mail address: `linda.keen@lehman.cuny.edu`