

## Discrete Structures: Sample Questions, Exam 1, Solutions

1. Prove by mathematical induction that  $3 \mid (n^3 - n)$  for every positive integer  $n$ .

**Answer:** Basis step  $n = 1$ :  $n^3 - n = 1^3 - 1 = 0$ , and so we must check that  $3 \mid 0$ . This is true since  $0 = 3(0)$ .

Induction step. For  $k \geq 1$ , assume  $P(k)$ , which is that  $3 \mid (k^3 - k)$ . We want to show  $P(k + 1)$ . In other words, we want to show that  $3 \mid [(k + 1)^3 - (k + 1)]$ .

So by  $P(k)$ , we know that there is an integer  $s$  satisfying

$$P(k) : \quad k^3 - k = 3s.$$

To show  $P(k + 1)$ , compute

$$\begin{aligned} [(k + 1)^3 - (k + 1)] &= (k^3 + 3k^2 + 3k + 1) - (k + 1) \\ &= k^3 + 3k^2 + 2k \\ &= (k^3 - k) + k + (3k^2 + 2k) \\ &= 3s + 3(k^2 + k) && \text{by } P(k) \\ &= 3(s + k^2 + k). \end{aligned}$$

Therefore  $3 \mid [(k + 1)^3 - (k + 1)]$ , and we have shown  $P(k + 1)$ .

Note to get from the second line to the third in the computation, we want to use  $P(k)$ . So therefore, in the second line, we want to find the expression  $k^3 - k$ . This is the motivation in going from the second to the third lines.

Since we have the basis step  $n = 1$  and the induction step, the proof is complete.

2. Write down a formula for the sequence

$$3, 4, 6, 9, 13, 18, 24, 31, 38, \dots$$

Is your formula recursive or explicit?

**Answer:**  $a_1 = 3$ .  $a_n = a_{n-1} + (n - 1)$ . This is a recursive formula.

3. Consider the matrices

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad \mathbf{B} = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \\ 1 & 0 & 1 \end{bmatrix}.$$

Compute  $\mathbf{A} \odot \mathbf{B}$ ,  $\mathbf{AB}$ ,  $\mathbf{B} \odot \mathbf{A}$  and  $\mathbf{A} \wedge \mathbf{B}$ .

Assume  $\mathbf{A}$  is the matrix of a relation. Draw the corresponding digraph.

**Answer:**


$$\mathbf{A} \odot \mathbf{B} = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 0 & 1 \end{bmatrix}$$

$$\mathbf{AB} = \begin{bmatrix} 1 & 1 & 2 \\ 1 & 1 & 1 \\ 1 & 0 & 1 \end{bmatrix}$$

$$\mathbf{B} \odot \mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 0 & 1 \end{bmatrix}$$

$$\mathbf{A} \wedge \mathbf{B} = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Digraph of  $\mathbf{A}$



4. Use the Euclidean algorithm to compute the greatest common divisor  $\text{GCD}(75, 12)$ . Show your work. Compute the least common multiple  $\text{LCM}(75, 12)$ .

**Answer:**

$$75 = 6(12) + 3$$

$$12 = 4(3) + 0$$

So the last one before the 0 is 3. So  $\text{GCD}(75, 12) = 3$ .

$$\text{LCM}(75, 12) = \frac{(75)(12)}{\text{GCD}(75, 12)} = \frac{(75)(12)}{3} = (75)4 = 300.$$

5. How many ways can a committee of 3 faculty members and 2 students be selected from 7 faculty members and 8 students? Show your work.

**Answer:** Task  $T_1$ : to choose 3 faculty members from 7, there are

$${}^7C_3 = \frac{7 \cdot 6 \cdot 5}{3 \cdot 2 \cdot 1} = 35$$

ways. Task  $T_2$ : to choose 2 students from 8, there are

$${}^8C_2 = \frac{8 \cdot 7}{2 \cdot 1} = 28$$

ways. All together, there are  $35 \cdot 28 = 980$  ways of choosing this committee.

6. Two fair six-sided dice are rolled and the sum  $s$  of the numbers coming up is recorded. What is the probability that  $s \geq 10$ ? Show your work.

**Answer:** The sample space

$$A = \{(1, 1), (1, 2), \dots, (6, 6)\},$$

and  $|A| = 36$ . The event that the sum is  $\geq 10$  is

$$E = \{(4, 6), (5, 5), (5, 6), (6, 4), (6, 5), (6, 6)\}.$$

So the probability is

$$\frac{|E|}{|A|} = \frac{6}{36} = \frac{1}{6}.$$

7. True/False. Circle T or F. No explanation needed.

- (a) T F If  $\mathbf{A}$  and  $\mathbf{B}$  are any  $2 \times 2$  matrices, then  $\mathbf{AB} = \mathbf{BA}$ .  
**Answer:** F—Matrix multiplication is not in general commutative.
- (b) T F If  $\mathbf{A}$ ,  $\mathbf{B}$  and  $\mathbf{C}$  are  $2 \times 2$  matrices, then  $(\mathbf{AB})\mathbf{C} = \mathbf{A}(\mathbf{BC})$ .  
**Answer:** T—Matrix multiplication is associative.
- (i) T F Let  $R = (A, \$)$  be the mathematical structure where  $A$  is the set of even integers and  $\$$  is a unary operation on  $A$  given by  $\$a = \frac{a}{2}$  for every  $a \in A$ .  $A$  is closed with respect to the operation  $\$$ .  
**Answer:** F—For  $a = 2$ ,  $\$a = 1 \notin A$ .
- (m) T F 49 and 77 are relatively prime.  
**Answer:** F— $\text{GCD}(49, 77) = 7 \neq 1$ .
- (n) T F Let  $A$  and  $B$  be subsets of a universal set  $U$ . Then it is always true that  $|A \cup B| = |A| + |B|$ .  
**Answer:** F—In general  $|A \cup B| = |A| + |B| - |A \cap B|$ , and so the equation is false if  $A$  and  $B$  are not disjoint.
- (o) T F Two cards are dealt in succession from a standard shuffled 52-card deck. The number of possible 2-card hands is 1326.  
**Answer:** T—The number of such hands is

$${}_{52}C_2 = \frac{52 \cdot 51}{2 \cdot 1} = 1326.$$

8. Let  $a_n$  be the sequence recursively defined by  $a_1 = 2$ ,  $a_2 = 3$ ,  $a_n = a_{n-1}a_{n-2}$  for  $n \geq 3$ . Compute the first five terms  $a_1, \dots, a_5$ .

**Answer:**

$$\begin{aligned} a_1 &= 2 \\ a_2 &= 3 \\ a_3 &= a_2a_1 = 2(3) = 6 && \text{(use } n = 3) \\ a_4 &= a_3a_2 = 6(3) = 18 \\ a_5 &= a_4a_3 = 18(6) = 108 \end{aligned}$$

9. A 6-sided die is rolled twice. What is the probability that the sum of the two rolls is exactly 8?

**Answer:** The sample space for two rolls of a die is

$$A = \{(1, 1), (1, 2), \dots, (6, 5), (6, 6)\},$$

and  $|A| = 6^2 = 36$ . The event given by the sum of the two rolls being 8 is given by

$$E = \{(2, 6), (3, 5), (4, 4), (5, 3), (6, 2)\},$$

and so  $|E| = 5$ . So the probability is

$$\frac{|E|}{|A|} = \frac{5}{36}.$$

10. Compute the truth table for the statement  $[(p \wedge q) \vee r] \Rightarrow (\sim q)$ . Show your work.

**Answer:**



$p$	$q$	$r$	$p \wedge q$	$(p \wedge q) \vee r$	$\sim q$	$[(p \wedge q) \vee r] \Rightarrow (\sim q)$
$T$	$T$	$T$	$T$	$T$	$F$	$F$
$T$	$T$	$F$	$T$	$T$	$F$	$F$
$T$	$F$	$T$	$F$	$T$	$T$	$T$
$T$	$F$	$F$	$F$	$F$	$T$	$T$
$F$	$T$	$T$	$F$	$T$	$F$	$F$
$F$	$T$	$F$	$F$	$F$	$F$	$T$
$F$	$F$	$T$	$F$	$T$	$T$	$T$
$F$	$F$	$F$	$F$	$F$	$T$	$T$

11. Consider the relation  $R$  on  $A = \{1, 2, 3, 4\}$  given by

$$1R2, \quad 2R3, \quad 3R3, \quad 3R4, \quad 4R3.$$

Draw the digraph of  $R$  and compute its connectivity relation  $R^\infty$ . Draw the digraph of  $R^\infty$ . (Hint: Try to determine  $R^\infty$  by inspection, not by computing with formulas.)

**Answer:**



Explanation for the extra edges in the digraph of  $R^\infty$ :

- $1R^\infty 3$  since  $1R2, 2R3$ .
- $1R^\infty 4$  since  $1R2, 2R3, 3R4$ .
- $2R^\infty 4$  since  $2R3, 3R4$ .
- $4R^\infty 4$  since  $4R3, 3R4$ .